Early Therapeutic Discovery in Oncology and Beyond: Challenges and Opportunities

Anton Simeonov, Ph.D.


Scientific Director, National Center for Advancing Translational Sciences (NCATS), National Institutes of Health (NIH)

TRACO Lecture September 30, 2019



Therapeutic discovery and development

Therapeutic discovery and development: the simplistic view and the reality



Translational inefficiency


The common causes of translational inefficiency are NCATS' focus

- Predicting safety and effectiveness of new drugs
- Scalable approaches to the >6000 untreatable diseases
- Data interoperability
- Biomarker qualification process
- Clinical trial networks
- Patient recruitment
- Electronic Health Records for research

- Harmonized IRBs
- Clinical diagnostic criteria
- Clinical outcome criteria (e.g., PROs)
- Adaptive clinical trial designs
- Shortening time of intervention adoption
- Methods to better measure impact on health (or lack thereof)
- Cross-sector collaborative structures
- Translational education/workforce development




High throughput screening



HTS is a standard step in the drug discovery process but has remained problem-ridden.

Quantitative high throughput screening

Improving the Process of Early Discovery: Quantitative High-Throughput Screening (qHTS)



PNAS 103:11473

- Conventional screening done at one concentration
 - Not appropriate for potency testing "dose makes the poison"
- qHTS tests compounds assayed at multiple concentrations (range: 4 logs)
- Enabled by miniaturized assay volumes (2-8 μL per test) and informatics pipeline
 - Generates pharmacological actives instead of statistical "hits"
 - Dramatically increases reliability
 - Dramatically reduces false positives and false negatives

 To date, several hundred million datapoints from several hundred screens have been generated and deposited in the public domain.

Full integration of medicinal chemistry

Full integration of medicinal chemistry to make more effective and less toxic tool molecules and drug candidates



Multi-parameter optimization (potency, solubility, metabolic stability)

~200 high-quality small molecule probes/tools to study novel therapeutic concepts and to shine light on the "dark matter of the genome" have been developed and made publicly available. Some of these are being further developed as drugs.

Small molecules

Example: small molecule probes to study cancer metabolism...with additional applications

- Tumors exhibit unique nutritional requirements. For example, pyruvate kinase M2 (PKM2) has been implicated in the Warburg Effect.
- A screen of PKM2, followed by medicinal chemistry optimization, produced several distinct molecules able to significantly activate the enzyme.
- These tool molecules have been used by multiple labs worldwide to study cancer biology and a subset are being further developed as therapies.



Inhibition of Pyruvate Kinase M2 by Reactive Oxygen Species Contributes to Cellular Antioxidant Responses Dimitrios Anastasiou ef al. Science 334, 1278 (2011); DOI: 10.1126/science.1211485

Ecumor IPrbs are sufficiently general and because density dependence and confirmmental variation affect most populations, those conclusions are Medy in retend to other systems. The construction and analysis of IPrbs across among of systems may provide support for this proposition, for addition to providing a test to suphers cross-evolutionary dynamics, IPrbs have also been exceeded to include quality variation and to identify-ovolutionarity states states (are 27, 27), giving them potential in unity several subdisciplines of population biology, including population gonetics, and

Inhibition of Pyruvate Kinase M2 by Reactive Oxygen Species Contributes to Cellular Antioxidant Responses

Disribrios Essetados, ^{1,4} Eistoge Poulogiaeris, ^{1,5} John M. Bussa, ^{1,5} Matthree M. Bussa, ⁴ Hanikang Jiang, ⁶ Min Shen, ⁶ Garp Bellinger, ^{1,5} Atsen T. Sassit, ^{1,5} Jason W. Boossile, ^{1,5} Douglas S. Auld, ⁶ Craig J. Thomas, ⁶ Marthew G. Vander Helden, ^{6,6} Lewis C. Castley^{6,5}†

A recent development: PKM2 activation may be a way to modulate metabolism in pancreatic beta cells to manage diabetes:

ARTICLES

medicine

Pyruvate kinase M2 activation may protect against the progression of diabetic glomerular pathology and mitochondrial dysfunction



Irreversible inhibition

SCIENCE TRANSLATIONAL MEDICINE | RESEARCH ARTICLE


CANCER

Irreversible inhibition of cytosolic thioredoxin reductase 1 as a mechanistic basis for anticancer therapy

William C. Stafford, ^{1,2} Xiaoxiao Peng, ¹* Maria Hägg Olofsson, ³† Xiaonan Zhang, ³ Diane K. Luci, ⁴ Li Lu, ⁵ Qing Cheng, ¹ Lionel Trésaugues, ⁶† Thomas S. Dexheimer, ⁴§ Nathan P. Coussens, ⁴¶ Martin Augsten, Hanna-Stina Martinsson Ahlzén, ^{1††} Owe Orwar, ^{2,7} Arne Östman, ^{3,8} Sharon Stone-Elander, ^{5,10} David J. Maloney, ⁴† Ajit Jadhav, ⁴ Anton Simeonov, ⁴ Stig Linder, ^{3,11} Elias S. J. Arnér ¹⁶⁶

- Thioredoxin reductase (TrxR): an underappreciated cancer target involved in control of redox balance.
- TrxR1 inhibitor, discovered from a robotic screen, acts by a covalent mechanism, yet is devoid of broad reactivity against similar targets (glutathione reductase, others).
- Unique selenocysteine residue in the enzyme active site responsible for the inhibitor's reactivity (i.e., covalent action highly context-dependent).
- The inhibitor is presently being developed by a new company spun out of Karolinska Institutet.

Copyright © 2018
The Authors, some rights reserved; exclusive licensee
American Association for the Advancement of Science, No claim to original U.S.
Government Works



E-mg/kg TRi-1 Group Day's Day's

FaDu xenograft, [18F]-FDG uptake





Example

Example: deep profiling of resistance development leads to an effective countermeasure

- Adaptive resistance in FMS-like receptor tyrosine kinase (FLT3)-mutant acute myeloid leukemia (AML) develops upon treatment with FLT3 inhibitors.
- To tackle the problem, the team examined the integrative in-cell kinase and gene regulatory network responses in tumors treated with FLT3 inhibitor.
- Innate immune pathway activation via the interleukin-1 receptor-associated kinase 1 and 4 (IRAK1/4) complex was found to contribute to adaptive resistance in FLT3mutant AML cells.

Targeting cancer metastasis

The other side of the "dark genome": targeting cancer metastasis as a translational challenge

Translational problem: 90% of cancer deaths are the consequence of metastatic spread, yet there are no drugs labeled as antimetastatic agents

The mechanisms of metastatic transformation and spread are not well elucidated

⇒ how do you develop therapies if you don't know the targets?

PNC & Metastasis

The Perinucleolar Compartment (PNC) is a nuclear organelle that appears in transformed cancer cells.

PNC prevalence correlates with breast cancer, colorectal cancer, and ovarian cancer disease progression, reaching nearly 100% in distant metastases.

PNCs are composed of a complex of proteins and RNA ⇒ no obvious single drug target

Approach: phenotypic screen for PNC disrupters as a therapeutic approach to block metastasis

Collaborative approach

Disease expert: Medicinal Chemistry: Clinical expert: Sui Huang, Northwestern University Jeffrey Aube, University of Kansas Udo Rudloff, NCI







Metarrestin

Discovery of Metarrestin

Through an unbiased phenotypic screen, the team identified a class of compounds that disrupt PNC in a metastatic tumor cell line (PC3-M) without cytotoxic effects.

A chemically-optimized molecule (metarrestin) demonstrated benefit in a mouse orthotopic model of pancreatic cancer metastasis.





Sci Transl Med. 2018 May 16;10(441)




Combination screening

Combination screening to overcome drug resistance in bacteria

Collaborators: Wei Zheng (NCATS), Peter Williamson (NIAID), Karen Frank (NIH Clinical Center)

- Developed assay and used qHTS to rapidly screen 5,170 approved drugs and drug candidates against multidrug-resistant (MDR) organisms.
- Tested thousands of two-drug combinations, identified synergistic pairs.





Metarrestin program


Metarrestin Program: Status and Next Steps

- Metarrestin is in pre-clinical development to target metastasis in pancreatic cancer.
- Pre-clinical development is currently in final stages through NCATS BrIDGs program (GMP manufacture & IND-enabling toxicity studies).
- Clinical support through NCI, goal is an IND filing in late 2019.




Drug combinations

Translation Challenge: Rapid Discovery of Drug Combinations



Resistance

Application of Drug Combinations to Address Resistance






Drug resistance


Dissemination of technology: combination screening to overcome drug resistance in cancer cells

- ABC subtype of Diffuse Large B-Cell Lymphoma (ABC-DLBCL) has poor prognosis and response to treatment
- Ibrutinib is a BTK inhibitor that has activity against ABC DLBCL
- Study evaluated 459 drugs in combination with Ibrutinib
 - » 6 x 6 concentration-response "matrix blocks", validation in 10 x 10 concentration-response matrix blocks
- DNA-damaging agents identified as synergizing with Ibrutinib in killing ABC DLBCL cell lines
- Dissemination:
 - » Protocols
 - Source code for dispense

High-throughput combinatorial screening identifies drugs that cooperate with ibrutinib to kill activated B-cell-like diffuse large B-cell lymphoma cells

Lodey A. Mathem Grinath", Rejechi Gebri", Ireal Shimh", Bush M. Youngh", Jonethan M. Koller", Songbo Liuf, Ian S. Geldest", Adem Yanger", Orpital Rickinight", Matthew B. Boser", Demies Y. Ouwest, Res-Rang Reng", San Michael", Yan Miscryesi', Werwel Heang", Martin I. Walsh', Bryan Y. Marti', Parsens Brosi"-, William Leister", Devid J. Malloney', Christopher A. Lodeir', Genesha Rei", Ajit Jadhey', Brian D. Payson', Christopher P. Austin', Sort E. Martin', Anton Simococki, Blazz Ferrer', Louis M. Staudh'', and Chief, J. Thomas''








Three-drug combinations

Three-drug combinations identified as effective against 10 common MDR clinical isolates



Sun W, et al. Emerging Microbes & Infections (2016)



3D models

Increasing the predictivity of in vitro assays: a continuum of 3D models of healthy and diseased tissues

2D Spheroids Organoids Printed Tissues Organ-on-a-chip

Physiological complexity


HTS compatibility

Physiological complexity



Tissue bioprinting

3D Tissue Bioprinting




Printed construct

The printer "3D prints" the cell/gel mixture in a layer by layer approach. The printed construct is incubated to allow the cells to form a tissue, and to enable proper cell differentiation.

Bioprinting

3D Bioprinting Projects





Native Blood Vessel Wall



Blood vessel wall (Kan Cao, U of Maryland)

Retina (Kapil Bharti, National Eye Institute)






Skin (Angela Christiano, Columbia University)



Epidermis

Layers of the Epidermis: native skin versus 3Dbioprinted skin




3D-Bioprinted Skin Stratum corneu **Epidermis** Stratum granulosum Stratum spino Stratum germinativu



Functional activity analyses

High-throughput tissue clearing protocols for highcontent, image and functional activity analyses


Clearing can be used to improve 3D tissue visualization and analysis



Stem cell technologies


Enabling Advanced 3D Models and Regenerative Medicine through Stem Cell Technologies

NCATS Stem Cell Translation Laboratory:




Access to relevant human cell types

Sensory neurons (nociceptors) and other neuronal subtypes, hepatocytes, etc.




Advanced imaging technologies for functional cell characterization

High-content confocal, calcium imaging, optogenetics



High-throughput electrophysiology methods

High-density multielectrode arrays 26,400 electrodes/well




Measurement of signaling pathways, metabolism & specific targets

Cyclic AMP, PKA activity, CREB phosphorylation, energy metabolism



Longitudinal tracking of cell behavior

Multiple measurements over days, weeks or months



Combined singlecell transcriptomic & proteomic analyses


Drug response in individual cells



Toxicity

Improved prediction of toxicity is crucial in drug development and environmental protection


Drug Development Failure Rates



Preclinical (21%) + Clinical (12%) Tox = 33% of all failures

Kola and Landis, Nature Reviews Drug Discovery 3, 711-716, 2004.

- There are over 80,000 chemicals in commerce, the majority with little to no toxicological data.
- We cannot solve the problem using laboratory animal tests only.



Collins, Gray, Bucher (2008) Science 319:906-7

Tox21 screening



The Tox21 Screening Project




Collection of diverse chemicals

In vitro test methods, screening High quality bioactivity data Predictive models
(of bioactivity of a new chemical in vitro and, one day, in vivo)



>50 screening campaigns of the 10K Collection

Tox21 10K Chemical Collection: ~10,000 chemicals (nominated and procured by EPA, NIEHS, and NCATS) comprising approved drugs, failed drugs, pesticides, industrial chemicals, etc. Extensive Quality Control →





Screening questions

Tox21 Screening Outcomes


- Rapid testing of chemicals enabled through robotic screening, largest collection of environmental chemicals and drugs assembled, multiple QC measures in place.
- Deposition into the public domain of the largest-ever toxicology dataset (almost 100M datapoints).
- Using crowdsourcing to move from data to knowledge.
- Estrogen receptor in vitro data being used by the EPA for regulatory purposes (https://www.epa.gov/endocrine-disruption/use-high-throughput-assays-and-computational-tools-endocrine-disruptor).
- Multiple organizations and consortia worldwide using Tox21 data (e.g., eTox/IMI, Tiley et al. Environ Int 101:19-26 used Tox21 data to rank chemicals of concern at Superfund sites).





Information sharing

Sharing internal know-how: Assay Guidance Manual (47 chapters/ 1,338 printed pages)






Table of Contents Considerations for Early Phase Drug Discovery 1 Chapter In Vitro Biochemical Assays 10 Chapters In Vitro Cell Based Assays 19 Chapters In VIvo Assay Guidelines 2 Chapters Assay Artifacts and Interferences 4 Chapters Assay Validation, Operations and Quality Control 5 Chapters Assay Technologies 2 Chapters Instrumentation 2 Chapters Pharmacokinetics and Drug Metabolism 1 Chapter Glossary of Quantitative Biology Terms 1 Chapter

Website: https://ncats.nih.gov/expertise/preclinical/agm

Email us: NCATS_AGM_Editors@mail.nih.gov



Linked in

Linkedin: www.linkedin.com/groups/7437344

https://ncats.nih.gov/agm-video

August 7th Videos

- Austin, CP: Welcome to the Assay Guidance Manual (AGM) Workshop.
- Coussers, NP: Strategies for Assay Selection & Robust Biochemical Assays
- Riss, T: Treating Cells as Reagents to Design Reproducible Screening Assays
- 4. Trask, OJ: Assay Development Considerations for High Content Imaging
- Auld, DS: Studies in Mechanisms and Methods in Assay Interferences
- 6. Dahlin, JL: Assay Interference by Chemical Reactivity
- Chung, TDY: Basic Assay Statistics, Data Analysis & Rules of Thumb
 - Devanarayan, V: Reproducibility & Differentiability of Potency Results
- Sittempelam, GS: Avoiding Artifacts & Interferences in Assay Operations

March 26-27th Videos

- Austin, CP: Welcome to the Assay Guidance Manual (AGM) Workshop
- 2. Coussens, NP: Robust Assays Define Success in Preclinical Research
- Lai-Nag, M: Target Identification & Validation in Translational Discovery
- 4. Foley, TL: Development & Validation of Cell-Based and Biochemical
- Riss, T: Treating Cells as Reagents to Design Reproducible Screening Assays
- Trask, OJ: Assay Development for HCS & Best Practices for 3D HCS.
- Roth, KD: Mass Spectrometry for Drug Screening and Lead Optimization
- 8. Dahlin, JL: Blossasy Interference by Aggregation and Chemical Reactivity
- 9. Patnalk, St Lead Selection and Optimization by Medicinal Chemistry
- 10. Xia, M: In Vitro Toxicological Testing Using a gHTS Platform
- 11. Xu, X: In Vitro Assessment of ADME Properties of Lead Compounds
- 12. Kahl, SD: Statistical Design of Experiments for Assay Development
- 13. Guha, R: Pheros Application to Target Evaluation and Drug Discovery
- Weidner, JR: Assay Operations: Keeping Assays Robust and Reproducible




Automating chemistry

Looking Ahead...Automating Chemistry as the Next Challenge in Preclinical Translation

Changes in biomedical research fields in the last 100 years:


Biology 1917








2017







Mapping biologically active chemical space to accelerate drug discovery

G. Sitta Sittampolom*, Dobrila D. Rudnicki, Danlio A. Tagle, Anton Simeonov and Christopher P. Austin

A specialized platform for innovative research exploration — ASPIRE — in preclinical drug discovery could help study unexplored biologically active chemical space through integrating automated synthetic chemistry, high-throughput biology and artificial intelligence technologies.

NATURE REVIEWS | DRUG DISCOVERY

doi:10.1038/d41573-018-00007-2

Published online 27 Nov 2018

Learn More About NCATS



Website: www.ncats.nih.gov



Facebook: facebook.com/ncats.nih.gov



Twitter: twitter.com/ncats_nih_gov



YouTube: youtube.com/user/ncatsmedia



E-Newsletter: ncats.nih.gov/news-andevents/e-news/e-news.html



Email us: info@ncats.nih.gov